

## Vector operations

### Representation of vectors

In the following, vector  $\mathbf{a}$  is represented  $\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$

with components in 3 directions east, north and up. Vectors are printed bold. In handwritten text, vectors are underlined or overscored with an arrow:  $\underline{\mathbf{a}}$   $\overline{\mathbf{a}}$

Magnitude of vector  $\mathbf{a}$  is  $a = |\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$

Unit vector in same direction  $\hat{\mathbf{a}} = \begin{pmatrix} a_1/a \\ a_2/a \\ a_3/a \end{pmatrix}$

### Basic vector operations

Vector addition

$$\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + b_3 \end{pmatrix}$$

Vector dot product

$$\mathbf{a} \cdot \mathbf{b} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \cdot \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = a_1 b_1 + a_2 b_2 + a_3 b_3$$

Vector dot product is a scalar  $\mathbf{a} \cdot \mathbf{b} = ab \cos \theta$  where  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$

Vector cross product

$$\mathbf{a} \times \mathbf{b} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \times \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{pmatrix} a_2 b_3 - b_2 a_3 \\ a_3 b_1 - b_3 a_1 \\ a_1 b_2 - b_1 a_2 \end{pmatrix}$$

Vector cross product is a vector with magnitude  $ab \sin \theta$  where  $\theta$  is the angle between  $\mathbf{a}$  and  $\mathbf{b}$ .  $\mathbf{a} \times \mathbf{b}$  is perpendicular to  $\mathbf{a}$  and  $\mathbf{b}$ . Vectors  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{a} \times \mathbf{b}$  form a right-handed set.

## Basic matrix algebra

Transpose of a matrix

$$\mathbf{A}^T = \begin{pmatrix} a & b \\ c & d \end{pmatrix}^T = \begin{pmatrix} a & c \\ b & d \end{pmatrix}$$

Symmetric matrix

$$\mathbf{A} = \begin{pmatrix} a & b & c \\ b & d & e \\ c & e & f \end{pmatrix} \text{ or } \mathbf{A} = \begin{pmatrix} a & b \\ b & d \end{pmatrix} \text{ or } a_{ij} = a_{ji}$$

Skew symmetric matrix

$$\mathbf{A} = \begin{pmatrix} a & b & c \\ -b & d & e \\ -c & -e & f \end{pmatrix} \text{ or } \mathbf{A} = \begin{pmatrix} a & b \\ -b & d \end{pmatrix} \text{ or } a_{ij} = -a_{ji}$$

Matrix addition (shown for 2 x 2 matrix)

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} + \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} a_{11} + b_{11} & a_{12} + b_{12} \\ a_{21} + b_{21} & a_{22} + b_{22} \end{pmatrix}$$

or in general  $a_{ij} + b_{ij} = c_{ij}$

Matrix multiplication (shown for 2 x 2 matrix)

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} = \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{pmatrix}$$

or, in general  $\sum_{p=0}^{p=n} a_{ip} b_{pj} = c_{ij}$

where  $n$  = number of dimensions

The unit matrix or Kronecker's  $\delta$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \text{ or } \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ or } \delta_{ij} \text{ or } \delta$$

Determinant of a 2 x 2 matrix

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Determinant of a 3 x 3 matrix

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = a \begin{vmatrix} e & f \\ h & i \end{vmatrix} - b \begin{vmatrix} d & f \\ g & i \end{vmatrix} + c \begin{vmatrix} d & e \\ g & h \end{vmatrix}$$

Eigenvectors of a matrix  $\mathbf{A}$  are solutions  $\mathbf{x}$  to:

$$\mathbf{A} \cdot \mathbf{x} = k \mathbf{x}$$

where  $k$  is an eigenvalue associated with eigenvector  $\mathbf{x}$ . For a symmetric matrix the eigenvectors are mutually perpendicular.